1-28-2019

Patient Perceptions of Planned Organ Removal During Hysterectomy

Zeinab Kassem
Chad M. Coleman
Andrew S. Bossick
Wan-Ting Su
Roopina Sangha
Ganesa Wegienka

Follow this and additional works at: <https://aah.org/jpcrr>

Part of the [Clinical Epidemiology Commons](#), [Medical Anatomy Commons](#), [Obstetrics and Gynecology Commons](#), [Public Health Education and Promotion Commons](#), [Surgery Commons](#), and the [Women's Health Commons](#)

Recommended Citation

Kassem Z, Coleman CM, Bossick AS, Su WT, Sangha R, Wegienka G. Patient perceptions of planned organ removal during hysterectomy. J Patient Cent Res Rev. 2019;6:28-35. doi: [10.17294/2330-0698.1658](https://doi.org/10.17294/2330-0698.1658)

Published quarterly by Midwest-based health system Advocate Aurora Health and indexed in PubMed Central, the Journal of Patient-Centered Research and Reviews (JPCRR) is an open access, peer-reviewed medical journal focused on disseminating scholarly works devoted to improving patient-centered care practices, health outcomes, and the patient experience.

Patient Perceptions of Planned Organ Removal During Hysterectomy

Zeinab Kassem, MD,¹ Chad M. Coleman, MPH,¹ Andrew S. Bossick, MPH,¹ Wan-Ting Su, PhD,¹ Roopina Sangha, MD, MPH,² Ganesa Wegienka, PhD¹

¹Department of Public Health Sciences, Henry Ford Health System, Detroit, MI; ²Department of Obstetrics and Gynecology, Henry Ford Hospital, Detroit, MI

Purpose	Previous reports indicate many women may not have a firm grasp on likely outcomes of different hysterectomy procedures. This study aimed to assess women's self-reported expectations of how they think their anatomy will change after hysterectomy.
Methods	Women scheduled for hysterectomy at a tertiary care hospital, for non-oncological reasons, reported their planned procedure type and the organs they understood would be removed 2 weeks prior to surgery. Patient reports and electronic medical records were reviewed, and kappa statistics (κ) were calculated to assess agreement for all women and within subgroups.
Results	Most of the 456 study participants (mean age: 48.02 ± 8.29 years) were either white/Caucasian ($n=238$, 52.2%) or African American ($n=196$, 43.0%). Among the 145 participants who reported a partial hysterectomy, 130 (89.7%) women indicated that their uterus would be removed and 52 (35.9%) reported that their cervix would be removed. Of those whose response was total hysterectomy ($n=228$), 208 (91.2%) participants reported their uterus would be removed and 143 (62.7%) reported their cervix would be removed. Among 144 women reporting a planned partial hysterectomy, only 15 (10.4%, $\kappa=0.05$) had a partial hysterectomy recorded in the electronic medical record. Among the 228 women who reported a planned total hysterectomy, 6.1% ($\kappa=0.05$) had a different procedure. While 125 participants reported planned ovary removal, only 93 (74.4%, $\kappa=0.55$) had an oophorectomy. Similarly, 290 participants reported planned fallopian tube removal, with 276 (95.2%, $\kappa=0.06$) having a salpingectomy.
Conclusions	A considerable proportion of women undergoing hysterectomy do not accurately report the organs that are planned be removed during their hysterectomy. This work demonstrates the need to improve patient understanding of their clinical care and its implications. (<i>J Patient Cent Res Rev</i> . 2019;6:28-35.)
Keywords	hysterectomy; patient education; patient expectations; gynecology; patient-centered outcomes

Since it was first performed in 1843,¹ hysterectomy has become the most common nonobstetrical surgery for women in the United States.^{2,3} Hysterectomy can involve removal of only the uterus (partial hysterectomy), removal of the uterus and the cervix (total hysterectomy), and removal of the uterus, cervix, and upper one-third to upper one-half of the vagina (radical hysterectomy). Hysterectomy may be

paired with additional surgeries, including removal of one or both ovaries (oophorectomy) or removal of the fallopian tubes (salpingectomy). The most common indication for hysterectomy is uterine fibroids,⁴ and most women report satisfaction and improvement in functioning after surgery.⁵⁻⁷ Due to the high frequency of hysterectomy, clinical research related to this surgery remains important.

Focus groups with 24 hysterectomy patients from an integrated health system based in Detroit, Michigan, indicated that some women were not able to report all procedures performed as part of their hysterectomy.⁸ More specifically, multiple women did not know if

Correspondence: Zeinab Kassem, MD,
Department of Public Health Sciences, 1 Ford Place, 5C,
Detroit, MI 48202 (kzeinab10@gmail.com)

they retained their ovaries, fallopian tubes, or cervix after their hysterectomy. Some women reported having a “total hysterectomy” and assumed that meant they no longer had ovaries or fallopian tubes. If women do not know whether they have these organs, they may not be receiving appropriate health care such as counseling regarding hormone replacement therapy and certain recommended screenings (eg, the Papanicolaou “Pap” test for cervical cancer screening). Women who cannot report whether they have ovaries may not be able to appropriately discuss their cardiovascular disease or cancer risk with health care providers.

The frequency of this lack of understanding is not well known. As this can have important health implications on the individuals involved, it is important to investigate in a systematic manner whether the results of the focus group study are isolated findings or whether deficiencies in understanding of hysterectomy is a more common problem that health care professionals should address. The aim of this work is to assess the accuracy with which patients report their expected anatomical changes from their hysterectomy and whether they varied by procedure indication or demographic characteristics.

METHODS

Study Population

Women scheduled for hysterectomy at Henry Ford Health System (Detroit, MI) who were at least 18 years of age and not undergoing surgery for suspected cancer or having a concomitant urogynecology procedure were invited to participate in a longitudinal cohort study to examine postsurgery regret. Eligible women were sent an introduction letter explaining the study and were contacted by telephone to complete a baseline interview in the 2 weeks before the patient’s scheduled surgery. Not all women who enrolled in the study had their planned surgery due to personal or clinical reasons. All women who completed the baseline questionnaires were included in the analyses of planned procedures and expected organ/tissue removal; however, only those who had a hysterectomy were included in analyses comparing self-report of surgical plans with actual surgical procedure performed.

The study protocols were approved by the local institutional review board, and consent was provided

by all participants. The study was funded through the Agency for Healthcare Research and Quality of the U.S. Department of Health and Human Services.

Data Collection

In the baseline questionnaire, women were asked which hysterectomy procedure they planned to have: total, partial, radical, or “don’t know.” They were then asked to select from the following list every organ/tissue they expected to be removed: ovary/ovaries, fallopian tube(s), partial/complete uterus, and partial/complete cervix. Answer options were presented in this manner to assess each woman’s understanding of clinical terminology. Women were asked to report their race and marital status (married or living as married vs not married or living as married). At 6 months posthysterectomy, the electronic medical record (EMR) of each participant was reviewed for surgical details, including surgical approach and organs/tissues removed as well as insurance type and indications for surgery, to assess accuracy of reporting within subgroups.

Statistical Analyses

Contingency tables, including frequencies and percentages, were created for the following pairs of variables: 1) the expected surgeries (partial/total hysterectomy) and the expected organs for removal; 2) the expected surgeries and the surgeries documented in the EMR; and 3) the EMR-documented surgeries and the reports of expected organ removal. Kappa coefficients with 95% CI were calculated to test the level of agreement between the two binary outcome variables.^{9,10}

RESULTS

Table 1 provides summary statistics for study participants. Mean age of the 456 participants was 48.0 ± 8.3 years, and most were either white/Caucasian ($n=238$, 52.2%) or African American ($n=196$, 43%). More than half the women were married or living as married and were employed full-time. Most of the participants (71.4%) had private insurance, and 27.1% had Medicaid and/or Medicare. Of the 456 women who enrolled, 452 (99.1%) women did undergo hysterectomy.

Expected Surgery vs Organ Removal

Among the 145 participants who reported that they were scheduled for a partial hysterectomy, 130 (89.7%)

Table 1. Descriptive Information About Participants Included in Analysis (N=456)

Characteristic	n (%)
Race	
African American	196 (43.0%)
White/Caucasian	238 (52.2%)
Others	22 (4.8%)
Current marital status	
Divorced	80 (17.5%)
Married/Living as married	244 (53.5%)
Never married	105 (23.0%)
Separated	10 (2.2%)
Widowed	17 (3.7%)
Employment status	
Employed full-time	254 (55.7%)
Employed part-time	46 (10.1%)
Self-employed	11 (2.4%)
Student and not working for pay	5 (1.1%)
Unemployed and not a student	110 (24.1%)
Others	30 (6.6%)
Insurance plan (N=462)*	
Private	330 (71.4%)
Medicare/Medicaid	125 (27.1%)
Military related	4 (0.9%)
No insurance	3 (0.6%)

**Some participants indicated they had more than 1 insurance type, thus insurance type is not mutually exclusive.*

indicated their uterus was going to be removed, 52 (35.9%) indicated that their cervix was going to be removed, and 11 (7.6%) reported that they were not sure which organs would be involved. Among the 228 participants who reported a plan for total hysterectomy, 208 (91.2%) reported that their uterus was going to be removed, 143 (62.7%) reported that their cervix was going to be removed, and 17 (7.5%) said that they were not sure which organs would be involved in the surgery. A total of 79 participants (17.3%) reported that they were not sure what type of surgery was planned. Among these 79 women unsure of the planned procedure type, 63 (79.8%) reported that their uterus was going to be removed and 27 (34.2%) reported that they expected removal of the cervix (Table 2).

After excluding women who reported “not sure” when indicating what surgery type they expected, 377 women remained in the analytical sample. Among those who reported they did not plan to have their cervix removed, only 93 (52%) women indicated they would be undergoing a partial hysterectomy (removal of the uterus with preservation of the cervix). Among those who expected removal of the cervix during surgery, 52 (26.3%) women reported they would be having a partial hysterectomy and 146 (73.7%) reported a different planned surgical approach. These results have a kappa coefficient of 0.26 (95% CI: 0.16–0.36), suggesting a low level of agreement between the expected surgery type and cervix removal (Table 3).

Among those who reported planned uterus and cervix removal, 142 (72.8%) women indicated they would be undergoing a total hysterectomy (removal of the uterus and cervix simultaneously). Among those who did not report planned removal of both the uterus and cervix during surgery, 86 (47.3%) women reported the planned surgery as total hysterectomy and 96 (52.7%) reported a different planned surgical approach. These results also had a kappa coefficient of 0.26 (95% CI: 0.16–0.36), suggesting participant reporting of type of surgery was not accurate for the reported planned organ removal (Table 3).

Expected Surgery vs Procedure Performed

Among the 144 women who reported a planned partial hysterectomy and whose EMR were reviewed, only 15 (10.4%) had a partial hysterectomy as recorded in their EMR. The agreement between participants’ reports of planned partial hysterectomy and official surgical reports was quite low, with a kappa coefficient of 0.05 (95% CI: -0.07–0.17). Among the 148 women who reported that they planned to not have a total hysterectomy, 133 (89.9%) participants did indeed have a total hysterectomy as recorded in their EMR ($\kappa=0.05$; 95% CI: -0.07–0.17) (Table 3).

Expected Organ Removal vs Procedure Performed

Among 37 participants with EMR-documented partial hysterectomy, 36 (97.3%) reported in the presurgery survey that their uterus was going to be removed and 13 (35.1%) reported that their cervix was going to be removed. For those with an EMR-

Table 2. Summary of Participant Self-Report of Planned Organ Removal and Self-Report of Planned Surgical Approach

Self-Report of Planned Organ Removal	Self-Report of Planned Surgical Approach				Total
	Partial Hysterectomy	Total Hysterectomy	Radical Hysterectomy	Not Sure	
	n=145	n=228	n=4	n=79	N=456
Uterus	130 (89.7%)	208 (91.2%)	3 (75.0%)	63 (79.8%)	404 (88.6%)
Cervix	52 (35.9%)	143 (62.7%)	3 (75.0%)	27 (34.2%)	225 (49.3%)

Percentages were calculated from respective column total.

documented total hysterectomy, 366 of 415 (88.2%) reported that their uterus was going to be removed and only 211 (50.8%) reported that their cervix was going to be removed. Among the 415 participants who underwent total hysterectomy, only 207 participants (49.9%) reported that both their uterus and cervix were to be removed. There was little agreement between expected and actual cervix removal ($\kappa=0.05$; 95% CI: -0.05–0.14) or expected and actual cervix and uterus removal ($\kappa=0.04$; 95% CI: -0.05–0.13). The agreement was similarly poor for salpingectomy ($\kappa=0.06$; 95% CI: -0.06–0.18). However, although only weak, the agreement was strongest for oophorectomy ($\kappa=0.55$; 95% CI: 0.47–0.64) (Table 3).

Overall, the kappa values did not vary by insurance status, age, marital status, race, or reason for hysterectomy, with the single exception of women who underwent hysterectomy because of endometriosis (data not shown). Among those having a hysterectomy due to endometriosis, the agreement between whether a woman expected to have her ovaries removed and whether she had them removed was strong ($\kappa=0.82$; 95% CI: 0.59–1.05).

DISCUSSION

While hysterectomy has been the most common nonobstetrical surgery in the United States for decades,^{2,3} it is not uncommon for women to inaccurately report the organs/tissues that will be removed during their planned hysterectomy. Prior work regarding this topic involving focus groups conducted by two separate research teams found that a significant number of women lack sufficient knowledge about the details of their hysterectomy and anatomy to

accurately report the details of their hysterectomy.^{8,11} Based on the information collected from participants in our study, in addition to the clinical data and surgical details provided by the EMR, the level of agreement between what patients expected and what physicians actually do in the operating room was tested. Most notably, nearly 10% of women who were planning to have a hysterectomy did not report that their complete or partial uterus would be removed and 36% of the women who reported a planned partial hysterectomy incorrectly thought that they would also have their cervix removed.

Similar results were reported in a cohort of 190 predominately white and college-educated women who had undergone a minimally invasive hysterectomy — only 67% were able to report correctly whether they had their cervix removed and whether they needed future cervical cancer screening.¹² Further, only 57% of the women understood the purpose of a Pap test.¹² Other publications also have demonstrated the lack of understanding of the female anatomy. In a study of college students that used a word bank to label a diagram of the female reproductive tract, less than 60% of the 138 women were able to correctly label the cervix and nearly 60% were able to identify the uterus.¹³ In a group of 500 patients (age 18–59 years) from an academic obstetrics and gynecology practice in the United States, 29% reported that hysterectomy included removal of the ovaries and tubes and approximately 25% of the women thought menstruation would continue after hysterectomy.¹⁴ The same study found 59% of the women did not report that removing the entire uterus would eliminate the possibility for cervical cancer, while two-thirds of the

Table 3. Comparison of Pre- and Postsurgical Data With Respect to Reported Planned Organ Removal, Surgical Approach, and Electronic Medical Record (EMR)

Self-Report of Planned Surgical Approach				
Self-Report of Planned Organ Removal	Partial Hysterectomy ^d	Not Partial Hysterectomy	Total N ^a	Kappa (95% CI)
Not cervix	93 (52%)	86 (48%)	179	0.26 (0.16–0.36)
Cervix	52 (26.2%)	146 (73.7%)	198	
Total	145	232	377	
	Total Hysterectomy ^d	Not Total Hysterectomy	Total N ^a	Kappa (95% CI)
Uterus + cervix	142 (72.8%)	53 (27.2%)	195	0.26 (0.16–0.36)
Not [uterus + cervix]	86 (47.3%)	96 (52.8%)	182	
Total	228	149	377	
EMR Report of Completed Surgical Approach				
Self-Report of Planned Surgical Approach	Partial Hysterectomy ^d	Not Partial Hysterectomy	Total N ^b	Kappa (95% CI)
Partial hysterectomy	15 (10.4%)	129 (89.6%)	144	0.05 (-0.07–0.17)
Not partial hysterectomy	14 (6.0%)	218 (94%)	232	
Total	29	347	376	
	Total Hysterectomy ^d	Not Total Hysterectomy	Total N ^b	Kappa (95% CI)
Total hysterectomy	214 (93.9%)	14 (6.1%)	228	0.05 (-0.07–0.17)
Not total hysterectomy	133 (89.9%)	15 (10.1%)	148	
Total	347	29	376	
EMR Report of Completed Surgical Approach				
Self-Report of Planned Organ Removal	Partial Hysterectomy ^d	Not Partial Hysterectomy	Total N ^c	Kappa (95% CI)
Not cervix	24 (10.5%)	204 (89.5%)	228	0.05 (-0.05–0.14)
Cervix	13 (5.8%)	211 (94.2%)	224	
Total	37	415	452	
	Total Hysterectomy ^d	Not Total Hysterectomy	Total N ^c	Kappa (95% CI)
[Uterus + cervix]	207 (94.1%)	13 (5.9%)	220	0.04 (-0.05–0.13)
Not [uterus + cervix]	208 (89.7%)	24 (10.3%)	232	
Total	415	37	452	
	Salpingectomy ^d	Not Salpingectomy	Total N ^c	Kappa (95% CI)
Yes	276 (95.2%)	14 (4.8%)	290	0.06 (-0.06–0.18)
No	146 (90.1%)	16 (9.9%)	162	
Total	422	30	452	
	Oophorectomy ^d	Not Oophorectomy	Total N ^c	Kappa (95% CI)
Yes	93 (74.4%)	32 (25.6%)	125	0.55 (0.47–0.64)
No	53 (16.2%)	274 (83.8%)	327	
Total	146	306	452	

^aTotal N: 377 (456 patients who had presurgery surveys – 79 self-reporting “not sure”).

^bTotal N: 376 (452 women who underwent hysterectomy – 76 (self-reporting “not sure”).

^cTotal N: 452 (patients who underwent hysterectomy).

^dPartial Hysterectomy = removal of uterus; Total Hysterectomy = removal of uterus + cervix; Salpingectomy = removal of fallopian tubes; Oophorectomy = removal of ovaries.

Percentages were calculated from respective row total.

women thought they would need to continue to have Pap tests after a total hysterectomy.¹⁴ Some of these results are not directly comparable to those reported herein, as the women in our analyses were actually planning to have a hysterectomy and had already completed presurgical counseling.

The findings suggest there should be concern about these women obtaining optimal future care. Surgical details are routinely collected as part of any health history, and care plans are affected by a patient's history. Giving inaccurate details about a health history or incomplete details about prior surgeries could lead to missed screening recommendations or inaccurate diagnoses or health score calculations. Reid et al found that women in the waiting room area at a women's health clinic had significant knowledge deficits regarding female pelvic anatomy and wanted their physicians to provide them with better education and counseling.¹⁵ Such findings highlight that already-existing systems of patient education are failing to properly educate women.

In a study by Wade et al, in which focus groups were conducted with hysterectomy patients in the 2-year period after their surgery, women expressed concern regarding lack of information and lack of participation in decision-making regarding their hysterectomy.¹¹ They reported feeling the quality and quantity of information about the surgery and their own anatomy and physiology was insufficient, specifically stating they wanted to know more about organ functions and which organs remained after surgery.¹¹ Women expressed a need to have their physicians spend more time with them and a need to have access to educational media. These issues are not unique to women undergoing hysterectomy.

Almost 20 years after the Wade et al¹¹ publication, Bossick et al⁸ conducted focus groups with women who had a hysterectomy at Henry Ford Health System. The women expressed similar concerns as those women 18 years prior. Not all women were able to report the type of hysterectomy they had or whether they had their tubes or ovaries removed, findings that spurred our current analysis. Further, our results along with information in the literature suggest that the lack of understanding appears to cross age and race groups.

These results highlight a persistent gap in women's understanding of hysterectomy. The reasons behind this knowledge gap could be due to lack of patient understanding of basic anatomy or challenges in surgical counseling and in-clinic patient education, but is likely a combination of these factors. Indeed, a 2013 study found that health literacy affects many U.S. women and plays an important role in women's health.¹⁶ In fact, health literacy's impact is not limited to women's health and affects health outcomes of other diseases like diabetes,¹⁷ inflammatory bowel disease,¹⁸ and heart failure.¹⁹ Lower health literacy also has been associated with worse health outcomes and quality of life in patients undergoing day surgery²⁰ as well as lower patient satisfaction and higher health care cost.²¹ Although our findings do not isolate the cause of the knowledge deficiency, the analyses do highlight an important opportunity to improve the care experience for women undergoing hysterectomy.

Based on the systematic review from Fredericks et al, and with media resources readily available, educational materials could be prepared and widely disseminated to better educate women about their own anatomy and how it will be affected by hysterectomy.²² Surgeons must counsel patients both before and after surgery to carefully explain any changes during surgery that deviated from the presurgical plan. This change in "routine care" must be led by clinicians who cannot assume anything about how well a patient understands her own anatomy. Indeed, Williams et al have shown that improving patient education on hysterectomy would improve patients' experiences.²³

Patient input should be sought for intervention development. For example, in a randomized control trial, women who were planning to undergo hysterectomy were randomized to usual care (n=52) or a scripted video (n=57) for their presurgical counseling.²⁴ The script for the video was created by the investigators and does not appear to have contained patient input. The participants were then asked true/false and multiple-choice questions about key information including knowledge about anatomy and the hysterectomy procedure. Those who were in the video arm had better knowledge scores initially, but not at 6 weeks after surgery.²⁴

Limitations

As with any epidemiological study, there are limitations to this work. The participant-reported information was collected within 2 weeks prior to hysterectomy and after presurgical counseling. During this time, surgical plans for some participants might have changed and this could have been communicated to the patient by their physician and could explain some, but likely not all, of the differences between planned hysterectomy type and the type that was performed. If the surgery had been explained to the woman in layman's terms, such as use of the word "womb" instead of uterus, we would not have captured this difference.

CONCLUSIONS

In summary, this work demonstrates the need to improve patient understanding of their clinical care and its implications. Hysterectomy is one of the most common surgeries for women, yet many women fail to understand how their body will change after this major surgery. A great opportunity exists to improve health literacy and educate women about their own basic anatomy, with a greater goal of improving their health and care satisfaction.

Patient-Friendly Recap

- Hysterectomy is a very common surgery and there are different types of hysterectomies that involve removal of different organs.
- The authors asked women undergoing hysterectomy to report the type of hysterectomy procedure planned and which organs would be removed. Surgical data from participants' electronic medical records also was collected.
- Many women were not able to accurately report the changes to their anatomy from the planned procedure.
- This work highlights the need to improve patient understanding of their clinical care, basic anatomy, and how both may change following surgery.

Author Contributions

Study design: Sangha, Wegienka. Data acquisition or analysis: all authors. Manuscript drafting: all authors. Critical revision: Kassem, Sangha, Wegienka.

Conflicts of Interest

None.

Funding Sources

This research was funded by the Agency for Healthcare Research and Quality (grant no. 5R24HS022417-05).

References

1. Sutton C. Hysterectomy: a historical perspective. *Baillieres Clin Obstet Gynaecol.* 1997;11:1-22. [CrossRef](#)
2. Torpy JM, Lynn C, Glass RM. JAMA patient page. Hysterectomy. *JAMA.* 2004;291:1526. [CrossRef](#)
3. Wilcox LS, Koonin LM, Pokras R, Strauss LT, Xia Z, Peterson HB. Hysterectomy in the United States, 1988-1990. *Obstet Gynecol.* 1994;83:549-55. [CrossRef](#)
4. Merrill RM. Hysterectomy surveillance in the United States, 1997 through 2005. *Med Sci Monit.* 2008;14(1):CR24-31.
5. Kuppermann M, Summitt RL Jr, Varner RE, et al. Sexual functioning after total compared with supracervical hysterectomy: a randomized trial. *Obstet Gynecol.* 2005;105:1309-18. [CrossRef](#)
6. Kuppermann M, Learman LA, Schembri M, et al. Predictors of hysterectomy use and satisfaction. *Obstet Gynecol.* 2010;115:543-51. [CrossRef](#)
7. Greer WJ, Richter HE, Wheeler TL, et al. Long-term outcomes of the Total or Supracervical Hysterectomy (TOSH) trial. *Female Pelvic Med Reconstr Surg.* 2010;16:49-57. [CrossRef](#)
8. Bossick AS, Sangha R, Olden H, Alexander GL, Wegienka G. Identifying what matters to hysterectomy patients: postsurgery perceptions, beliefs, and experiences. *J Patient Cent Res Rev.* 2018;5:167-75. [CrossRef](#)
9. McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med (Zagreb).* 2012;22:276-82. [CrossRef](#)
10. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med.* 2005;37:360-3.
11. Wade J, Pletsch PK, Morgan SW, Menting SA. Hysterectomy: what do women need and want to know? *J Obstet Gynecol Neonatal Nurs.* 2000;29:33-42. [CrossRef](#)
12. Mattingly M, Juran R, Su I, Ebinger J, Daggy J, Tucker Edmonds B. Patient knowledge of hysterectomy and pap screening after minimally invasive hysterectomy. *Patient Educ Couns.* 2017;100:121-5. [CrossRef](#)
13. Volck W, Ventress ZA, Herbenick D, Hillard PJ, Huppert JS. Gynecologic knowledge is low in college men and women. *J Pediatr Adolesc Gynecol.* 2013;26:161-6. [CrossRef](#)
14. Harmanli O, Ilarslan I, Kirupanathan S, Knee A, Harmanli A. Women's perceptions about female reproductive system: a survey from an academic obstetrics and gynecology practice. *Arch Gynecol Obstet.* 2014;289:1219-23. [CrossRef](#)
15. Reid JA, Templeman CL, Groneberg DA, Brueggmann D, Jaque JM. Patients' knowledge of female pelvic health and related educational preferences. *J Community Health.* 2017;42:147-54. [CrossRef](#)
16. Corrarino JE. Health literacy and women's health: challenges and opportunities. *J Midwifery Womens Health.* 2013;58:257-64. [CrossRef](#)
17. Saeed H, Saleem Z, Naeem R, Shahzadi I, Islam M. Impact of health literacy on diabetes outcomes: a cross-sectional study from Lahore, Pakistan. *Public Health.* 2018;156:8-14. [CrossRef](#)

18. Tormey LK, Reich J, Chen YS, et al. Limited health literacy is associated with worse patient-reported outcomes in inflammatory bowel disease. *Inflamm Bowel Dis*. 2019;25:204-12. [CrossRef](#)
19. Matsuoka S, Tsuchihashi-Makaya M, Kayane T, et al. Health literacy is independently associated with self-care behavior in patients with heart failure. *Patient Educ Couns*. 2016;99:1026-32. [CrossRef](#)
20. Hälleberg Nyman M, Nilsson U, Dahlberg K, Jaensson M. Association between functional health literacy and postoperative recovery, health care contacts, and health-related quality of life among patients undergoing day surgery: secondary analysis of a randomized clinical trial. *JAMA Surg*. 2018;153:738-45. [CrossRef](#)
21. MacLeod S, Musich S, Gulyas S, et al. The impact of inadequate health literacy on patient satisfaction, healthcare utilization, and expenditures among older adults. *Geriatr Nurs*. 2017;38:334-41. [CrossRef](#)
22. Fredericks S, Guruge S, Sidani S, Wan T. Postoperative patient education: a systematic review. *Clin Nurs Res*. 2010;19:144-64. [CrossRef](#)
23. Williams RD, Clark AJ. A qualitative study of women's hysterectomy experience. *J Womens Health Gend Based Med*. 2000;9 Suppl 2:S15-25. [CrossRef](#)
24. Pallett AC, Nguyen BT, Klein NM, Phippen N, Miller CR, Barnett JC. A randomized controlled trial to determine whether a video presentation improves informed consent for hysterectomy. *Am J Obstet Gynecol*. 2018;219:277.e1-e7. [CrossRef](#)

© 2019 Aurora Health Care, Inc.